| | | 2021 Blue Cross Blue Shield of Michgian Cardiovascular Consortium | | |
|---|--------|---|---------------|--------------|
| Collaborative Quality Initiative Performance Index Scorecard PCI & VS Combined Measurement Period: 01/01/2021 - 12/31/2021 | | | | |
| Measure # | Weight | Measure Description | PCI Points | VS Points |
| 1 | 10 | Meeting Participation - Clinician Lead | | |
| | | 2 Meetings (Vascular Surgery); 3 Meetings (PCI) | 5 | 5 |
| | | 1 Meeting (Vascular Surgery); 2 Meetings (PCI) | 2.5 | 2.5 |
| | | Did not participate | 0 | 0 |
| 2 | 5 | Data Coordinator Expectations (Vascular Surgery: Includes 1 year follow-up ≥80%) | | |
| | | Meets all expectations (1 Year FU <u>></u> 80%) | 2.5 | 2.5 |
| | | Meets most expectations (1 Year FU 60-79%) | 1 | 1 |
| | | Does not meet expectations (1 Year FU <60%) | 0 | 0 |
| 3 | 5 | Internal Case Reviews | | |
| | | Submitted reviews for ≥90% of cases | 2.5 | 2.5 |
| | | Submitted reviews for <90% of cases | 0 | 0 |
| 4 | 5 | Physicians Complete Cross Site Review of Assigned Cases for Procedural Indications and Technical Quality | | |
| | | Submitted reviews for 100% of cases | 2.5 | 2.5 |
| | | Submitted reviews for <100% of cases | 0 | 0 |
| 5 | 12.5 | Vascular Surgery Collaborative Goal - Statin at Discharge for Open Bypass, CEA* and CAS^ Discharges | | |
| | | ≥95% | NA | 12.5 |
| | | 93% - <95% | NA | 10 |
| | | 90% - <93% | NA | 5 |
| | | <90% | NA | 0 |
| 6 | 10 | Vascular Surgery Performance Goal – Surgeons to prescribe a maximum of 10 opioid pills for opioid naïve patients with EVAR ⁺ at discharge | | |
| | | ≥80% | NA | 10 |
| | | 75% - <80% | NA | 7.5 |
| | | 70% - <75% | NA | 5 |
| | | <70% | NA | 0 |
| 7 | 10 | Vascular Surgery Performance Goal – Surgeons to prescribe a maximum of 10 opioid pills for opioid naïve patients with CEA* at discharge | | |
| | | ≥80% | NA | 10 |
| | | 75% - <80% | NA | 7.5 |
| | | 70% - <75% | NA | 5 |
| | | <70% | NA | 0 |
| 8 | 12.5 | PCI Performance Goal - Peak Intra-Procedure ACT [‡] recorded | | |
| | | ≥90% | 12.5 | NA |
| | | 80% - <90% | 10 | NA |
| | | 70% - <80% | 5 | NA |
| | | <70% | 0 | NA |
| 9 | 10 | PCI Performance Goal - Percent of cases with peak ACT‡ ≥350 seconds for Heparin- only cases | | |
| | | ≤15% | 10 | NA |
| | | >15 - 25% | 7.5 | NA |
| | | >25% - 35% | 5 | NA |
| | | >35% | 0 | NA |
| 10 | 10 | NEW - PCI Performance Goal - Percent of cases with Air Kerma dose S Gray | | |
| | | <u><</u> 2% | 10 | NA |
| | | >2% - 3% | 7.5 | NA |
| | | >3% - 4% | 5 | NA |
| 11 | 10 | >4% NEW - PCI Collaborative Goal - Pre PCI hydration (oral and/or IV) (volume/3ML/Kg) in patients with eGFR** < 60 (excludes dialysis, cardiac arrest, cardiogenic shock, PCI status of "salvage" and symptomatic heart failure NYHA^A 2,3,4, and STEMI ⁺⁺). | 0 | NA |
| | | ≥50% | 10 | NA |
| | | 40% - <50% | 5 | NA |
| | | <40% | 0 | |
| | 100 | 10/07 | 0 | NA |

¹⁰⁰

28 sites participate in PCI & Vascular Surgery *CAS=carotid artery stent ^CEA=carotid endarterectomy *EVAR=endovascular aneurysm repair ‡ACT=activated clotting time **eGFR=estmated glomerular filtration rate

^^NYHA= New York Heart Association heart failure class

++STEMI=ST elevated myocardial infarction

Indicates Participation Measures

Indicates Performance Measures